

Research article

Open Access

Rapid evolution in response to introduced predators II: the contribution of adaptive plasticity

Leigh C Latta IV¹, Jeremy W Bakelar¹, Roland A Knapp² and Michael E Pfrender*¹

Address: ¹Department of Biology, 5305 Old Main Hill Road, Utah State University, Logan, UT 84322, USA and ²Sierra Nevada Aquatic Research Laboratory, University of California, HCR 79, Box 198, Mammoth Lakes, CA 93546, USA

Email: Leigh C Latta - leigh@biology.usu.edu; Jeremy W Bakelar - jwbakelar@cc.usu.edu; Roland A Knapp - knapp@lifesci.ucsb.edu; Michael E Pfrender* - pfrender@biology.usu.edu

* Corresponding author

Published: 14 February 2007

Received: 9 September 2006

BMC Evolutionary Biology 2007, **7**:21 doi:10.1186/1471-2148-7-21

Accepted: 14 February 2007

This article is available from: <http://www.biomedcentral.com/1471-2148/7/21>

© 2007 Latta et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Introductions of non-native species can significantly alter the selective environment for populations of native species, which can respond through phenotypic plasticity or genetic adaptation. We examined phenotypic and genetic responses of *Daphnia* populations to recent introductions of non-native fish to assess the relative roles of phenotypic plasticity versus genetic change in causing the observed patterns. The *Daphnia* community in alpine lakes throughout the Sierra Nevada of California (USA) is ideally suited for investigation of rapid adaptive evolution because there are multiple lakes with and without introduced fish predators. We conducted common-garden experiments involving presence or absence of chemical cues produced by fish and measured morphological and life-history traits in *Daphnia melanica* populations collected from lakes with contrasting fish stocking histories. The experiment allowed us to assess the degree of population differentiation due to fish predation and examine the contribution of adaptive plasticity in the response to predator introduction.

Results: Our results show reductions in egg number and body size of *D. melanica* in response to introduced fish. These phenotypic changes have a genetic basis but are partly due to a direct response to chemical cues from fish via adaptive phenotypic plasticity. Body size showed the largest phenotypic change, on the order of nine phenotypic standard deviations, with approximately 11% of the change explained by adaptive plasticity. Both evolutionary and plastic changes in body size and egg number occurred but no changes in the timing of reproduction were observed.

Conclusion: Native *Daphnia* populations exposed to chemical cues produced by salmonid fish predators display adaptive plasticity for body size and fecundity. The magnitude of adaptive plasticity was insufficient to explain the total phenotypic change, so the realized change in phenotypic means in populations exposed to introduced fish may be the result of a combination of initial plasticity and subsequent genetic adaptation. Our results suggest that immediately following the introduction of fish predators, adaptive plasticity may reduce the impact of selection through "Baldwin/Bogert effects" by facilitating the movement of populations toward new fitness optima. Our study of the response of a native species to an introduced predator enhances our understanding of the conditions necessary for rapid adaptive evolution and the relationship between rapid evolution and adaptive phenotypic plasticity.

Background

Introductions of non-native species can result in strong selective challenges for native populations. The strength of selection in this case is determined by the size of the environmental shift, which imposes a fitness cost on the population proportional to the squared distance between the population mean phenotype and the position of the new optimum [1,2]. If the optimum moves far enough, the fitness cost will be sufficiently high to reduce the intrinsic rate of increase of the population to <1 . Unless the population can rapidly advance toward the new optimum phenotype, it will not persist [3]. Two processes can facilitate persistence of populations challenged with a rapidly changing environment: adaptive phenotypic plasticity and genetic adaptation.

Adaptive phenotypic plasticity allows individuals within a population to accommodate a changing environment [4,5] by facilitating rapid movement to a new fitness optimum. This movement occurs through changes in the mean value of a trait and/or changes in the genetic and phenotypic variance/covariance structures. In the extreme case, plastic changes in the mean value of a trait are able to completely move a population to a new fitness optimum and no genetic adaptation is required [5]. In cases where a plastic change in the mean is not sufficient to shift a population to a new optimum it can allow a population to persist until sufficient adaptive genetic changes occur [6,7]. An incomplete shift in the population mean towards a new selective optimum that facilitates population persistence is called the "Baldwin effect" [6] and the resulting reduction in the intensity of selection is referred to as the "Bogert effect" [8] or adaptive buffering [9]. Plastic changes in the (co)variance matrix may result in increased levels of expressed genetic variance (*i.e.*, variance that is context-dependent and arises only in response to specific environmental cues) and changes in covariances between traits that increase the response to selection [10-13].

Populations may also adapt genetically to new environmental conditions when there is no pre-existing adaptive phenotypic plasticity or plasticity is insufficient to completely shift a population to a new phenotypic optimum. The rate of genetic adaptation toward a new optimum is determined by a number of factors, including the amount of additive genetic variation present for the traits under selection [14], the rate at which mutation produces new adaptive variation [15,16], and genetic correlations among characters [16-19].

A common source of rapid environmental change arises from the introduction of novel predator species. In a notable example, non-native fishes have been widely introduced into naturally fishless alpine lakes throughout the

world and have had profound effects on native zooplankton species, including *Daphnia*. *Daphnia* have a long history as a model system to study the consequences of introduced fish predators [20-23]. *Daphnia* adapt to introduced fish through changes in traits related to detection avoidance, including alterations in patterns of diel vertical migration (DVM) [24,25] and reduced body size [21,23,26]. *Daphnia* also display significant adaptive phenotypic plasticity in response to chemical cues produced by fish that can facilitate persistence during changes in selection regime. Plastic changes that reduce pigmentation [27] and body size [28-32] in *Daphnia* decrease the ability of fish to detect their prey resulting in higher survivorship, while plastic increases in fecundity [28-30] result in higher intrinsic rates of population increase.

Daphnia melanica (identified as *Daphnia middendorffiana* in previously published studies, *e.g.*, [33-35], but recently classified as *D. melanica* based on molecular analyses [M. Pfreder, unpublished data]) populations located in alpine lakes throughout the Sierra Nevada in eastern California, USA provide a unique opportunity to study the effects of introduced predators on naive populations. These alpine lakes have been the subjects of extensive ecological study [33-35] in part because the history of fish introductions is well documented. In lakes where *D. melanica* and fish co-occur, *D. melanica* have smaller body sizes and reproduce earlier relative to those in lakes without fish [36]. These differences were attributed to rapid adaptive evolution. However, because *Daphnia* are often highly plastic in response to chemical cues from fish, the differences in morphology and life-histories observed previously may not be solely due to underlying genetic alteration. Differences in morphology and life-history could be entirely due to phenotypic plasticity or a combination of plastic and genetic modification.

To determine the relative contributions of adaptive plasticity and genetic adaptation during rapid evolution in response to introduced fish we conducted common-garden experiments on clonally reproducing females of *D. melanica* populations collected from four lakes in the Sierra Nevada with contrasting fish stocking histories. Two lakes were never stocked and remain in their natural fishless condition and two lakes have contained introduced fish populations during the last several decades. We measured morphological and life-history traits of clonally reproducing females from each population in the presence and absence of chemical cues from fish (*i.e.*, fish kairomone). Because *D. melanica* can be maintained in a state of constant clonal reproduction in the lab, it is straightforward to utilize standard quantitative genetic techniques to estimate the contribution of genetic and plastic phenotypic effects underlying adaptive traits. Our chief working assumption in this experiment is that the

phenotypic states of fishless populations are representative of the ancestral phenotypic states of populations that currently contain fish. Given our assumption of equality between currently fishless populations and ancestral states of fish populations is true, our study design allowed us to determine the degree of morphological and life-history differentiation due to selection by fish predation and quantify the contribution of phenotypic plasticity in determining adaptive responses to the introduction of fish.

Results

Number of eggs and fecundity

Due to occasional mortality in the life-table prior to release of first clutch we measured egg number as an index calibration for fecundity to increase our sample sizes. For individuals that had both egg number and number of live offspring measured egg number was a highly significant predictor of the number of viable offspring (all regressions: $p < 0.01$). All regressions (described in more detail in the methods section) showed a positive correlation between the two variables. Correlation coefficients for the data subsets were between 0.43 and 0.57.

Levels of phenotypic plasticity

Fish kairomone caused significant reductions in mean body size at maturity for all populations (Fig. 1A). In the kairomone(-) treatment, mean body size at maturity for all genotypes was 1.78 mm, while the average size at maturity in the kairomone(+) treatment was 1.70 mm. Non-significant interaction terms suggest that a population's response to fish kairomone is independent of its history of fish introductions. Mean age at maturity did not change in response to fish kairomone (Fig 1B). This result appears largely as a consequence of the large variances associated with this trait. The number of eggs in the brood pouch increased in response to fish kairomone (Fig. 1C). The number of eggs increased significantly from 4.1 in kairomone(-) to 5.2 in kairomone(+) in response to fish kairomone (Table 1). Although there is a tendency for fishless populations to produce more eggs in response to kairomones than fish populations, the difference in reaction norms between fishless and fish populations is not significant so changes in clutch size are also independent of the history of fish introductions.

Because levels of variation for a trait are often context dependent we calculated coefficients of variation for body size, age at maturity, and egg number to determine if the amount of variance in these traits is dependent upon the presence/absence of fish kairomone. Coefficients of variation were lowest for body size at maturity (range 1.5 – 9.8), intermediate for age at maturity (range 10.5 – 22.1), and highest for egg number (range 32.8 – 51.4). Expressed variance showed little response to fish kairomone based

on our criterion of non-overlapping confidence intervals. Variance in body size increased significantly only in the Frog Lake population in the presence of fish kairomone (Table 2).

Genetic differentiation among populations

Genotypes from fishless populations had significantly larger body sizes at maturity than did genotypes from populations that co-exist with fish (Table 1). The average body size of genotypes from fishless lakes was 1.97 mm while genotypes from fish-containing lakes averaged 1.51 mm in size. Post hoc pairwise comparisons based on t values between all populations showed that genotypes from fishless populations ("Source" and "Frog") did not differ from one another ($p = 0.47$), but are significantly larger than genotypes from Puppet Lake ($p < 0.0001$) and Evelyn Lake ($p < 0.0001$). Puppet Lake genotypes are also significantly larger than genotypes from Evelyn Lake ($p < 0.0001$). A non-significant interaction term implies that these differences are not sensitive to the presence of fish kairomone (Table 1).

There was a significant reduction in the number of eggs in the brood pouch, from 5.27 eggs per individual in fishless populations to 4.05 eggs per individual in populations co-occurring with fish (Table 1). However, this result appears largely influenced by one population. Post hoc comparisons show that Evelyn Lake genotypes produce significantly fewer eggs than genotypes from Frog, Source, and Puppet Lake ($p = 0.0015$, 0.0298, and 0.0241, respectively), but Frog, Source, and Puppet Lakes do not differ in egg production (all possible pairs: $p > 0.2602$). A non-significant interaction suggests the difference in egg production between Evelyn Lake genotypes and all others did not depend on the assay environment. Age at maturity did not differ among fish and fishless populations (Table 1).

Discussion

Rapid evolution is an important component of the success of invading species [37] and the response of organisms in invaded communities [9] because it ameliorates the selective cost imposed by a shift in the phenotypic optimum and enhances the probability of long-term population persistence. Similarly, adaptive plasticity may be an important component of rapid evolution as it can allow short-term population persistence following changes in the selective environment [6,8] that in turn provides time for evolutionary mechanisms to operate. However, disentangling actual cases of rapid evolution from purely plastic changes in response to a changing environment can be problematic because adaptive phenotypic plasticity is a common feature in many organisms [38].

In this study we examined the rapid changes of *Daphnia* morphology and life histories in response to a single




Figure 1
Population responses to fish kairomone. Reaction norm plots for a) body size, b) age at maturity, and c) number of eggs in response to presence (+) or absence (-) of fish kairomone. Open symbols left of the vertical dotted lines are values for fish-less populations, filled symbols right of the vertical dotted lines are values for fish-containing populations. Error bars are +/- one standard error of the mean.

Table 1: NANOVA results.

Effect	Trait									
	df	Size			Age			Egg #		
		F	p		df	F	p	df	F	p
Environment	1/60	6.87	0.0111	1/60	0.81	0.3726	1/60	5.41	0.0235	
Type	1/60	278.56	<0.0001	1/60	0.42	0.5196	1/60	5.85	0.0186	
Pop(Type)	2/60	66.06	<0.0001	2/60	1.79	0.1760	2/60	3.11	0.0519	
Environment × Pop(Type)	2/60	0.02	0.9776	2/60	3.13	0.0507	2/60	0.12	0.8890	
Environment × Type	1/60	0.71	0.4014	1/60	0.00	0.9945	1/60	3.35	0.0723	

Results from NANOVA for morphological (size) and life-history (age and egg number) traits. Shown are the degrees of freedom (df), F-values (F) and p-values (p). Significant results (p < 0.05) are indicated in bold.

abrupt change in the environment. Specifically, we investigated the relative role of genetic and plastic phenotypic changes in moving populations exposed to a novel predator toward a new fitness optimum. Our results show that reductions in egg number and body size of *D. melanica* genotypes from the Sierra Nevada, in response to introduced fish, are largely adaptive evolutionary responses and not due entirely to adaptive phenotypic plasticity. We do find evidence for adaptive plasticity, in the form of increases in clutch size and reductions in body size, in these populations that could facilitate short-term persistence and subsequent rapid evolution. We caution, however, that our interpretation of these results is predicated on the assumption that our measured phenotypes of currently fishless populations are representative of the ancestral phenotypes of populations that currently co-exist with fish.

Our results suggest that naive *D. melanica* populations in the Sierra Nevada may initially respond to fish introductions through adaptive phenotypic plasticity brought about by chemical cues from fish, which facilitates movement towards the new phenotypic optimum. First, plastic reductions in body size make *D. melanica* less visible to fish and constitute evidence for a "Baldwin/Bogert effect" [6,8]. Fish are highly effective size-selective predators and their efficiency is primarily linked to prey visibility [26,39-41]. Thus, *D. melanica* that are less visible have a fitness advantage (via increased survivorship) through movement towards the new phenotypic optimum and the resulting reduction in selection intensity due to decreased predator efficiency. Concomitant with a decrease in body size, *D. melanica* also show adaptive phenotypic increases in fecundity that could lead to higher intrinsic rates of population increase. Although the rate at which fish

Table 2: Means and CV's.

Population	N	Exposure Time	Trait	Treatment			
				Kairomone (-)		Kairomone (+)	
				Mean (SE)	CV (95% CI)	Mean (SE)	CV (95% CI)
Source	7	0	Size	2.01 (0.04)	5.0 (3.2–10.9)	1.96 (0.04)	5.1 (3.3–11.3)
			Age	9.82 (0.50)	13.4 (8.6–30.3)	9.65 (0.38)	10.5 (6.7–23.4)
			Egg #	4.00 (0.72)	47.8 (28.9–165.2)	5.86 (0.74)	33.3 (20.8–86.9)
Frog	9	0	Size	1.98 (0.01)	1.5 (1.0–2.9)	1.93 (0.06)	9.8 (6.6–19.0)
			Age	10.18 (0.65)	19.2 (12.8–38.1)	9.63 (0.38)	11.7 (7.9–22.8)
			Egg #	4.44 (0.60)	40.8 (26.3–95.2)	6.78 (1.09)	48.2 (30.7–125.4)
Puppet	10	53	Size	1.77 (0.03)	6.2 (4.3–11.4)	1.68 (0.04)	7.7 (5.3–14.2)
			Age	10.94 (0.64)	18.6 (12.7–34.5)	9.21 (0.53)	18.2 (12.4–34.3)
			Egg #	4.60 (0.48)	32.8 (22.0–66.6)	5.10 (0.72)	44.7 (29.3–101.5)
Evelyn	8	91	Size	1.34 (0.03)	6.0 (3.9–12.2)	1.24 (0.02)	5.6 (3.7–11.5)
			Age	8.54 (0.46)	15.2 (10.0–31.8)	9.55 (0.75)	22.1 (21.5–47.6)
			Egg #	3.25 (0.59)	51.4 (31.8–163.2)	3.25 (0.45)	39.4 (25.0–99.1)

Estimates of phenotypic means and coefficients of variation. The units for body sizes are mm and for ages at maturity are days. Standard errors for the mean and 95% Modified McKay confidence intervals for coefficients of variation are given in parentheses.

remove individuals from these *Daphnia* populations is unknown, our observed increase in clutch size of approximately one is quite significant. Estimates of *D. melanica* population sizes in the Sierra Nevada are on the order of hundreds of millions to billions [R. Knapp, unpublished data], thus, an increase of one individual at first reproduction might substantially offset any losses due to predation. An interesting aspect of our findings is that our naive *Daphnia* populations, those without any history of fish exposure, are responsive to chemicals produced by fish. This observation suggests that *D. melanica* may be pre-adapted to fish predation, and that the genetic machinery responsible for adaptive phenotypic plasticity in response to fish kairomone is ancestral in this species.

Although we find evidence for adaptive phenotypic plasticity that would facilitate short-term population persistence in the face of novel predation, it is not sufficient to explain the difference in body-size and egg number between populations that are historically fishless and those that co-occur with fish. For example, a comparison of the average body size in the Evelyn Lake population (1.34 mm in the kairomone(-) and 1.24 mm in the kairomone(+) treatments) with the average in the fishless populations (2.00 mm in the kairomone(-) and 1.95 mm in the kairomone(+) treatments) shows that the mean phenotype in Evelyn Lake has diverged by 9.4 phenotypic standard deviations. The change in body size attributable to plasticity in fishless populations is approximately one standard deviation. In other words, the change in body size due to plasticity accounts for only about 11% of the total difference observed between Evelyn and fishless populations. Thus, the phenotypic differences observed in our study appear largely due to changes in the underlying genetic components controlling phenotype.

Our observation that the body-size response in Evelyn Lake was much higher than that in Puppet Lake could arise for three reasons. First, the difference in body size could simply reflect the different amounts of time each population was exposed to fish predation (Puppet Lake – 53 years; Evelyn Lake – 91 years). Second, *Daphnia* populations may have experienced differing levels of fish predation resulting in varying selection intensities, with the selection intensity in Evelyn Lake substantially higher. Finally, given our observation that expressed levels of genetic variance for body size increased approximately 6-fold in response to fish kairomone in one fishless population (Frog Lake) but not in the other (Source Lake) our fish populations may have differed in the initial levels of standing genetic variation, with the Evelyn Lake population harboring more standing genetic variation than Puppet Lake.

Our observation that Evelyn Lake was the only population to display a significant evolutionary reduction in egg number is likely due to the ability of *D. melanica* to deposit large amounts of melanin in the carapace. Melanin production in the carapace would initially "blind" selection to changes in egg number. Thus, an evolutionary response in egg number should occur only after reductions in melanin deposition. Fish predation produces strong selection on melanin production in other *Daphnia* populations [42], and there is evidence for reduced melanin expression in *Daphnia* from our fish populations relative to fishless populations [M. Pfreder, unpublished data]. Therefore, the apparent delayed onset of selection on egg number could be due to initial selection on melanin production and subsequent selection on egg number.

Traditional views of character evolution typically involve trade-offs among traits that can limit the adaptive potential of a population [43]. However, several selection experiments involving *Daphnia* suggest adaptive evolutionary changes in one trait are not necessarily associated with concomitant maladaptive changes in others [44-46]. We observe a similar result here, where evolutionary and plastic changes in body size and fecundity occur in the absence of changes in the timing of maturity and reproduction. Our results, and those of other researchers that imply the absence of a trade-off, could be attributed to assay conditions in which food is not a limiting resource [47]. *Daphnia* morphology and life-history can display food concentration-dependent reactions to the presence of fish kairomone [48].

Conclusion

We investigated the relative contributions of selection and adaptive phenotypic plasticity to the rapid evolution of morphology and life histories in response to an introduced predator. We conclude that adaptive plasticity could facilitate short-term population persistence through "Baldwin/Bogert effects", but that long-term persistence was achieved through subsequent genetic adaptation. Further investigation into other traits that may have also undergone rapid change in selective regime as a consequence of fish introductions, such as pigmentation and DVM behavior, examined under differing kairomone and food conditions, will provide a more detailed view of the traits and processes involved in the overall evolution of the *Daphnia*/fish predator-prey system in the Sierra Nevada.

Numerous studies have examined the contributions of plasticity and selection to rapid adaptation in non-native species following their introduction into a novel environment (*e.g.*, [49]). In contrast, few studies have examined the phenotypic and evolutionary response of native species to introduced species that pose strong novel selective

challenges. Thus, this study and a growing body of others investigating the response of native communities to introduced species should enhance our understanding of the conditions necessary for rapid adaptive evolution and the relationship between rapid evolution and population persistence [9].

Methods

Study populations

Individual genotypes used in the life-table assay were collected from four permanent lakes in the central Sierra Nevada during the summer of 2004. These lakes are located in the Humphreys, French Canyon, and Vogelsang basins at elevations ranging from 3150–3632 meters. Frog Lake (ID# 52103; UTM Zone 11: 351079 E, 4124432 N) and Source Lake (UTM Zone 11; 349988 E 4125708 N), remain in their natural fishless condition (referred to collectively as fishless populations). Puppert and Evelyn Lakes were naturally fishless but were stocked with trout during the past century. Puppert Lake has been stocked with golden trout (*Oncorhynchus mykiss aguabonita*) every other year since 1951 (California Dept. of Fish and Game, unpublished stocking records), resulting in 53 years of fish predation on the resident *D. melanica* population at the time of collection. Evelyn Lake (UTM Zone 11; 295393 E, 4186659 N) was initially stocked with brown trout (*Salmo trutta*) in 1913. Brook trout (*Salvelinus fontinalis*) were introduced in 1928, 1946, 1947, 1949, 1951, 1954 and 1958, and rainbow trout (*Oncorhynchus mykiss*) were introduced in 1939, 1942, 1944, 1957, 1962, and 1966 [50]. No stocking has occurred since 1966, and the resident rainbow trout population is self-sustaining. In total, *Daphnia* in Evelyn Lake were exposed to 91 years of fish predation at the time of collection (Puppert Lake and Evelyn Lake are referred to collectively as fish populations).

Clone establishment and maintenance

Daphnia were collected from each of the study lakes and maintained at 4°C for a period of 1–2 weeks prior to isolation in the lab. To capture the maximum amount of genetic variation from each population, mature females from the original field collection were isolated and placed singly in 250 mL beakers containing 200 mL of filtered well-water. This procedure ensures that no isolates were genotypically identical juveniles produced in the period from collection in the field until isolation in the lab. Isolated individuals were maintained by clonal reproduction under constant conditions of temperature (15°C) and 16L:8D photoperiod for approximately 20 generations prior to experimentation. Water levels in the beakers were kept constant with the periodic addition of double-distilled water. *Daphnia* were fed a vitamin supplemented pure culture of the green alga *Scenedesmus obliquus* every 3–4 days.

Life-table assay

Morphological and life-history characteristics were assayed using a standard experimental design [51,52]. Briefly, single immature females were taken from the stock isolates, each representing an experimental line. The lines were then maintained as single asexually produced progeny for two generations. In third generation individuals, we measured a suite of traits upon reaching maturity (defined as the first instar with the deposition of eggs into the brood pouch). Two traits, number of eggs in the brood pouch and size at maturity are directly related to visibility and potential for survival in the face of visually-feeding predators. The two remaining traits, age at maturity and number of viable offspring produced are related to the intrinsic rate of population increase. Each experimental line was maintained in a 250 mL beaker containing 150 mL of filtered well-water supplemented with a constant concentration (135,000 cells/mL) of *S. obliquus*. Upon reaching maturity, second generation lines assigned to the fish kairomone treatment were placed in filtered well-water aged with a 20–25 cm bull trout (*Salvelinus confluentus*) for 24 hours. (the kairomone treatment is referred to as kairomone(+)) and the non-kairomone treatment as kairomone(-)) Exposing second generation individuals to fish kairomone post-maturity ensures that maternal effects due to fish kairomone are minimized. All beakers in the life-table assay were maintained in a controlled temperature room with a 16L:8D photoperiod at 18°C and their position in the chamber changed every two days to minimize micro-environmental differences. The food/water mixture in all beakers was replaced with food/water of the appropriate type, kairomone(+) or kairomone(-), every other day.

Statistical procedures

We performed linear regression on egg number and number of surviving offspring upon release of first clutch to determine if egg number serves as a proxy for the more general fitness character of fecundity. Regressions were run on four separate subsets of the data: 1) fishless populations in the kairomone(-) treatment; 2) fishless populations in the kairomone(+) treatment; 3) fish populations in kairomone(-) treatment; and 4) fish populations in the kairomone(+) treatment. Analyzing the subsets separately aided in determining whether a correlation between egg number and viable offspring is sensitive to environmental and/or genetic differences between populations.

Nested analysis of variance (NANOVA) in which covariance parameters were estimated using restricted maximum likelihood was performed on three traits (body size at maturity, age at maturity, and egg number at maturity) to test for fixed effects of treatment (kairomone(+) or kairomone(-)), lake type (fish or fishless), population nested within lake type, and interactions between environment

and lake type, and between environment and population nested within lake type [PROC MIXED; [53]]. The model was designed to account for heterogeneity in covariance matrices across treatments because variance and covariance estimates can vary across environments.

Interpretation of results based on our model is relatively straightforward. A significant treatment effect is evidence for phenotypic plasticity in a given trait, irrespective of a populations' fish stocking history. A significant lake type effect implies phenotypic differences between populations in historically fishless and fish-containing lakes. The strength of conclusions about the actual level of genetic differentiation underlying phenotypic divergence is based on the level of significance of the interaction term. For example, a significant effect of lake type in conjunction with non-significant interaction terms would indicate underlying genetic differences among populations regardless of treatment effects.

Plasticity in the expressed genetic variance of traits was assessed by calculating coefficients of variation (CV) for each population across treatments separately. We then constructed 95% Modified McKay confidence intervals for each CV [54,55] and assessed differences between estimates based on the degree of overlap of confidence intervals.

Authors' contributions

LCL designed and conducted the experiment, performed statistical analyses, and drafted the manuscript. JWB conducted the experiment. RAK provided background information on *Daphnia* populations, assisted with fieldwork in the Sierra Nevada, and helped draft the manuscript. MEP conceived of the study, conducted fieldwork in the Sierra Nevada to collect the experimental populations, and contributed to analysis and manuscript preparation. All authors read and approved the final manuscript.

Acknowledgements

The authors wish to thank C. Werner for assistance in the laboratory. The Utah State University Evolution Group, S. Carroll, F. Messina, A. Scoville, and an anonymous reviewer provided helpful comments on the manuscript. Funding for this study was provided by NSF grant DEB-021212487, a USU New Faculty Research Grant, and a USU Center for Integrated BioSystems research grant to M. Pfrender, and NSF grant DEB-0075509 to R. Knapp and O. Sarnell.

References

- Lynch M, Lande R: **Evolution and extinction in response to environmental change.** In *Biotic Interactions and Global Change* Edited by: Kareiva PM, Kingsolver JG, Huey RB. Sunderland: Sinauer; 1993:234-250.
- Lande R, Shannon S: **The role of genetic variation in adaptation and population persistence in a changing environment.** *Evolution* 1996, **50**:434-437.
- Gomulkiewicz R, Holt RD: **When does evolution by natural selection prevent extinction?** *Evolution* 1995, **49**:201-207.
- Dudley SA, Schmitt J: **Testing the adaptive plasticity hypothesis: Density-dependent selection on manipulated stem length in *Impatiens capensis*.** *Am Nat* 1996, **147**:445-465.
- Price TD, Qvarnstrom A, Irwin DE: **The role of phenotypic plasticity in driving genetic evolution.** *Proc R Soc Lond B Biol Sci* 2003, **270**:1433-1440.
- Baldwin JM: **A new factor in evolution.** *Am Nat* 1896, **30**:441-451.
- Yeh PJ, Price TD: **Adaptive phenotypic plasticity and the successful colonization of a novel environment.** *Am Nat* 2004, **164**:531-542.
- Huey RB, Hertz PE, Sinervo B: **Behavioral drive versus behavioral inertia in evolution: a null model approach.** *Am Nat* 2003, **161**:357-366.
- Strauss SY, Lau JA, Carroll SP: **Evolutionary response of natives to introduced species: what do introductions tell us about natural communities?** *Ecol Lett* 2006, **9**:357-374.
- Schlichting CD: **The evolution of phenotypic plasticity in plants.** *Annu Rev Ecol Syst* 1986, **17**:667-693.
- Boersma M, Spaak P, De Meester L: **Predator-mediated plasticity in morphology, life history, and behavior of *Daphnia*: The uncoupling of responses.** *Am Nat* 1998, **152**:925-927.
- Pigliucci M, Cammell K, Schmitt J: **Evolution of phenotypic plasticity: a comparative approach in the phylogenetic neighbourhood of *Arabidopsis thaliana*.** *J Evol Biol* 1999, **12**:779-791.
- Pigliucci M: **Evolution of phenotypic plasticity: where are we going now?** *Trends Ecol Evol* 2005, **20**:481-486.
- Fisher RA: *The Genetical Theory of Natural Selection* New York: Dover; 1958.
- Falconer DS, Mackay TFC: *Introduction to Quantitative Genetics* Essex: Prentice Hall; 1996.
- Lynch M, Walsh B: *Genetics and Analysis of Quantitative Traits* Sunderland: Sinauer Associates; 1998.
- Arnold SJ: **Constraints on phenotypic evolution.** *Am Nat* 1992, **140**:S85-S107.
- Etterson JR: **Evolutionary potential of *Chamaecrista fasciculata* in relation to climate change. Genetic Architecture of three populations reciprocally transplanted along an environmental gradient in the great plains.** *Evolution* 2004, **58**:1459-1471.
- Etterson JR, Shaw RG: **Constraint to adaptive evolution in response to global warming.** *Science* 2001, **294**:151-154.
- Dodson SI: **Complementary feeding niches sustained by size-selective predation.** *Limnol Oceanogr* 1970, **15**:131-137.
- Wells L: **Effects of alewife predation on zooplankton populations in Lake Michigan.** *Limnol Oceanogr* 1970, **15**:556-565.
- Werner EE, Hall DJ: **Optimal foraging and the size selection of prey by bluegill sunfish (*Lepomis macrochirus*).** *Ecology* 1974, **55**:1042-1052.
- Kitchell JA, Kitchell JF: **Size-selective predation, light transmission, and oxygen stratification: Evidence from the recent sediments of manipulated lakes.** *Limnol Oceanogr* 1980, **25**:389-402.
- Pijanowska J, Weider LJ, Lampert W: **Predator-mediated genotypic shifts in a prey population: Experimental evidence.** *Oecologia* 1993, **96**:40-42.
- Cousyn C, De Meester L, Colbourne JK, Brendonck L, Verschuren D, Volckaert F: **Rapid, local adaptation of zooplankton behavior to changes in predation pressure in the absence of neutral genetic changes.** *Proc Natl Acad Sci USA* 2001, **98**:6256-6260.
- Galbraith MGJ: **Size-selective predation on *Daphnia* by rainbow trout and yellow perch.** *Trans Am Fish Soc* 1967, **96**:1-10.
- Tollrian R, Heibl C: **Phenotypic plasticity in pigmentation in *Daphnia* induced by UV radiation and fish kairomones.** *Funct Ecol* 2004, **18**:497-502.
- Stibor H: **Predator induced life-history shifts in a freshwater cladoceran.** *Oecologia* 1992, **92**:162-165.
- Reede T: **Life history shifts in response to different levels of fish kairomones in *Daphnia*.** *J Plankton Res* 2005, **17**:1661-1667.
- Reede T, Ringleberg J: **The influence of a fish exudates on two clones of the hybrid *Daphnia galeata* × *hyalina*.** *Cladocera as Model Organisms in Biology: Proceedings of the Third International Symposium On Cladocera* 1995, **307**:207-212.
- Spaak P, Vanoverbeke J, Boersma M: **Predator-induced life-history changes and the coexistence of five taxa in a *Daphnia* species complex.** *Oikos* 2000, **89**:164-174.

32. Sakwinska O: **Response to fish kairomone in *Daphnia galeata* life history trait relies on shift to earlier instar at maturation.** *Oecologia* 2002, **131**:409-417.
33. Bradford DF, Cooper SD, Jenkins TM Jr, Kratz K, Sarnelle O, Brown AD: **Influences of natural acidity and introduced fish on faunal assemblages in California alpine lakes.** *Can J Aquat Sci* 1998, **55**:2478-2491.
34. Knapp RA, Matthews KR, Sarnelle O: **Resistance and resilience of alpine lake faunal assemblages to fish introductions.** *Ecol Monogr* 2001, **71**:401-421.
35. Knapp RA, Hawkins CP, Ladau J, McClory JG: **Fauna of Yosemite National Park lakes has low resistance but high resilience to fish introductions.** *Ecol Appl* 2005, **15**:835-847.
36. Fisk DF, Latta LC IV, Knapp RA, Pfrender ME: **Rapid evolution in response to introduced predators I: rates and patterns of morphological and life-history trait divergence.** *BMC Evolutionary Biology* . in review
37. Lee CE: **Evolutionary genetics of invasive species.** *Trends Ecol Evol* 2002, **17**:386-391.
38. Pigliucci M: *Phenotypic plasticity: Beyond nature and nurture* Baltimore, MD: Johns Hopkins University Press; 2001.
39. Brooks JL, Dodson SI: **Predation, body size, and composition of plankton.** *Science* 1965, **150**:28-35.
40. Zaret TM, Kerfoot WC: **Fish predation on *Bosmina longirostris*, body size selection versus visibility selection.** *Ecology* 1975, **56**:232-237.
41. O'Brien WJ, Kettle D, Riessen H: **Helmets and invisible armor: structures reducing predation from tactile and visual planktivores.** *Ecology* 1979, **60**:287-294.
42. Saegrov H, Hobaek A, L'Abée-Lund JH: **Vulnerability of melanic *Daphnia* to brown trout predation.** *J Plankton Res* 1996, **18**:2113-2118.
43. Roff DA: *Life History Evolution* Sunderland, MA: Sinauer Associates; 2002.
44. Spitze K: **Chaoborus predation of life-history evolution in *Daphnia pulex*: temporal pattern of population diversity, fitness and mean life history.** *Evolution* 1991, **45**:82-95.
45. Spitze K, Burnson J, Lynch M: **The covariance structure of life-history characters in *Daphnia pulex*.** *Evolution* 1991, **45**:1081-1090.
46. Baer CF, Lynch M: **Correlated evolution of life-history with size at maturity in *Daphnia pulicaria*: patterns within and between populations.** *Genet Res* 2003, **81**:123-132.
47. Reznick D, Nunney L, Tessier A: **Big houses, big cars, superfleas and the costs of reproduction.** *Trends Ecol Evol* 2000, **15**:421-425.
48. Weetman D, Atkinson D: **Antipredator reaction norms for life-history traits in *Daphnia pulex*: dependence on temperature and food.** *Oikos* 2002, **98**:299-307.
49. Dybdahl MF, Kane SL: **Adaptation vs. phenotypic plasticity in the success of a clonal invader.** *Ecology* 2005, **86**:1592-1601.
50. Elliot MJ, Loughlin MH: **Historical overview of fishery management in Yosemite National Park 1877-1992.** In *Natural Resources Report* draft. Yosemite National Park, California; 2005.
51. Lynch M: **Spontaneous mutations for life-history characters in an obligate parthenogen.** *Evolution* 1985, **39**:804-818.
52. Pfrender ME, Lynch M: **Quantitative genetic variation in *Daphnia*: temporal changes in genetic architecture.** *Evolution* 2000, **54**:1502-1509.
53. SAS Institute Inc. Cary, NC. 2004.
54. McKay AT: **Distribution of the coefficient of variation and the extended 't' distribution.** *J Roy Stat Soc* 1932, **95**:695-698.
55. Vangel MG: **Confidence intervals for a normal coefficient of variation.** *American Statistician* 1996, **15**:21-26.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

