

Correction

Open Access

Quantitative genetic analysis of life-history traits of *Caenorhabditis elegans* in stressful environments

Simon C Harvey*, Alison Shorto and Mark E Viney

Address: School of Biological Sciences, University of Bristol, Woodland Road, Bristol, BS8 1UG, UK

Email: Simon C Harvey* - Simon.Harvey@bristol.ac.uk; Alison Shorto - Alison.Shorto@bristol.ac.uk; Mark E Viney - Mark.Viney@bristol.ac.uk

* Corresponding author

Published: 12 May 2009

Received: 11 May 2009

BMC Evolutionary Biology 2009, **9**:96 doi:10.1186/1471-2148-9-96

Accepted: 12 May 2009

This article is available from: <http://www.biomedcentral.com/1471-2148/9/96>

© 2009 Harvey et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

In Figure 1 of [Harvey et al (Evolutionary Biology 2008, 8:15)] the plotted data were inverted. The correct Figure is shown below. The text and statistical analyses in [Harvey et al (Evolutionary Biology 2008, 8:15)] are correct.

Correction

In Figure 1 of [1] the plotted data were inverted. The correct Figure is shown below. The text and statistical analyses in [1] are correct.

References

1. Harvey SC, Shorto A, Viney ME: **Quantitative genetic analysis of life-history traits of *Caenorhabditis elegans* in stressful environments.** *BMC Evolutionary Biology* 2008, **8**:15.




Figure 1

***C. elegans* varies in its dauer larvae formation phenotype.** (A) The proportion of dauer larvae that developed at two concentrations of food (1 and 2% w/v *E. coli* OP50 in water) for 163 RILs and (B) this for 21 RILs and N2 (■) and DR1350 (□) in one representative assay.