

Research article

Open Access

Shape differences rather than size differences between castes in the Neotropical swarm-founding wasp *Metapolybia docilis* (Hymenoptera: Vespidae, Epiponini)

Mário V Baio¹, Fernando B Noll*^{1,2} and Ronaldo Zucchi¹

Address: ¹Departamento de Biologia, Faculdade de Filosofia Ciências e Letras de Ribeirão Preto, Universidade de São Paulo, 14040-901-Ribeirão Preto (SP), Brazil and ²Museum of Biological Diversity, Department of Entomology, The Ohio State University, 1315 Kinnear Road, Columbus, OH 43212; USA

Email: Mário V Baio - mv@zipmail.com.br; Fernando B Noll* - fernandobn@ffclrp.usp.br; Ronaldo Zucchi - rzucchi@ffclrp.usp.br

* Corresponding author

Published: 13 May 2003

Received: 16 January 2003

BMC Evolutionary Biology 2003, 3:10

Accepted: 13 May 2003

This article is available from: <http://www.biomedcentral.com/1471-2148/3/10>

© 2003 Baio et al; licensee BioMed Central Ltd. This is an Open Access article: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

Abstract

Background: Swarm-founding epiponine wasps are an intriguing group of social insects in which colonies are polygynic (several queens share reproduction) and differentiation between castes is often not obvious. However, caste differences in some may be more pronounced in later phases of the colony cycle.

Results: Using morphometric analyses and multivariate statistics, it was found that caste differences in *Metapolybia docilis* are slight but more distinct in latter stages of the colony cycle.

Conclusions: Because differences in body parts are so slight, it is proposed that such variation may be due to differential growth rates of body parts rather than to queens being larger in size, similar to other previously observed epiponines.

Background

Polistine wasps in the tribe Epiponini are remarkable for their polygyny, swarm-founding habits, and reputed slight differentiation between castes. Although the overall features of epiponines are well-documented and discussed [1–3], many sociobiological details remain obscure. For example, for decades, swarm-founding polistines have been considered permanently polygynous, even though monogyny has been detected in some species [1,4,5]. An explanation was offered by [6], who showed that *Metapolybia aztecoides* alternates between polygyny and oligogyny and, eventually monogyny in the course of its life cycle. This cyclical oligogyny [6] seems to be important in the maintenance of high genetic relatedness in epiponine wasps [7–9]. In fact, new queens are produced with just one queen in the nest [10] in some species.

Preliminary overall studies [1,4] demonstrated a wide range of queen-worker dimorphism among different taxa in the Epiponini. Three different forms of queen-worker distinction were recognized [1], but, recently, five forms of caste differentiation due to the influence of the colony cycle were recognized [11]. They are: 1, the absence of morphological differences between queens and workers during the entire colony cycle (*Parachartergus smithii* [12], *Pseudopolybia vespiceps* [13], *Chartergellus communis* [14], *Brachygastra lecheguana* [15]); 2, the absence of morphological differences between castes, with some young females developing ovaries in only some phases of the colony cycle (*Synoeca cyanea* [16]); 3, morphological discontinuities between castes related to variation during the colony cycle, with non-inseminated laying females present during the entire colony cycle (*Protopolybia exigua*

and *P. sedula* [11]); 4, caste differences increase during the colony cycle with non-inseminated laying females occurring only some phases of the colony cycle (*Chartergus globiventris* [11], *Polybia scutellaris*, *P. occidentalis*, and *P. paulista* [17], *Protonectarina sylveirae* [11], *Epipona guerini* [18]) and; 5, queens always distinct from workers, with no variation in this pattern occurring during the colony cycle (*Agelaia areata* [19], *A. vicina* [20,21], *A. pallipes* and *A. multipicta* [22], *Apoica flavissima* [23] and *A. pallens* [24], *Polybia dimidiata* [25]).

Even though *Metapolybia* may be considered a model genus for studies related to colony population structure in epiponines, just a few studies (taxonomy – [26], ovary development – [27], colony composition, including caste differences [28] and nest construction – [29]) have been performed on this taxon since [6], despite the increase in knowledge of this tribe. This paper examines caste differences and other ecological aspects of *Metapolybia docilis*.

Results

Nests and related aspects

Colonies under investigation were assigned to one of 4 stages, after [17] (Table 1): (a) pre-emergence of workers, in which no adult offspring had been produced (one colony); (b) emergence of workers, with different-aged brood and at least one adult generation (one colony); and (c) male producing (one colony). Also, two other colonies were migrating for a new location. This fourth stage was defined as (d) swarm or absconding (two colonies). Even though these two forms of swarm are difficult to distin-

guish, behavioral observations (unpublished data) indicated that only part of the population migrated to a new nest in the *swarm* colony. In addition, swarming queens were exclusively younger females (see below) suggesting production of sexuals in the mother nest. The absconding colony represents the migration of a whole colony after some disturbance once the nest was damaged. In addition all queens were old (see below) evidencing that this colony was not producing new queens, representing the end of a cycle.

Females' ovary condition, insemination and their relative age

Two types of ovarian development were recognized as type A, with filamentous ovarioles bearing no visible or slightly developed oocytes, and type B, with well developed and longer ovarioles, each one bearing from two to several mature oocytes. Since only type B females contained sperm in the spermatheca, females with type A and B ovarian development can be characterized as workers and queens, respectively.

According to our data, the colony cycle in *Metapolybia docilis* is characterized by great age variation (Figure 1). In the pre-emergence colony both workers and queens varied substantially in age. In worker-producing and male-producing colonies most workers were younger and most queens were older. In the absconding colony, workers varied in age, but the single queen was old (black pattern). In the swarm colony, queens and workers were young and old individuals, respectively, without intermediate age.

Table 1: Number of combs, brood production, and colony population in the analyzed colonies of *Metapolybia docilis*. Percentage in parentheses.

Nest Location	Colony phase	Cells				total	Estimated n. of generations	Females		Males
		Empty	with eggs	With larvae	With pupae			Workers	Queens	
Igaçaba – São Paulo State	Worker-producing	123	108	268	321	820	1	197	23	0
		(15.0)	(13.2)	(32.7)	(39.2)			(89.6)	(10.5)	
Igaçaba – São Paulo State	Male-producing	313	42	26	6	387	2	165	4	7
		(79.9)	(10.9)	(6.7)	(1.6)			(93.8)	(2.3)	(3.98)
Pedregulho – São Paulo State	Pre-emergence	41	19	0	0	60	0	64	9	0
		(68.3)	(31.7)					(87.7)	(12.3)	
Nova Xavantina – Mato Grosso State	absconding	-	-	-	-	-	-	44	1	0
								(97.8)	(2.2)	
Ribeirão Preto – São Paulo State	Swarm	-	-	-	-	-	-	11	7	0
								(61.1)	(38.9)	




Figure 1
 Frequency of queens (White bars) and workers (dark bars) according to the color patterns of the transverse apodeme from the 4th gastral sternite in *Metapolybia docilis* in different analyzed colonies. LY – light yellow, LB – light brown, DB – dark brown, B – black.

Morphological differences

Mean differences of 10 measured characters were tested in workers and queens by Bonferroni *t*-test (Table 2). In all analyzed colonies queens were hardly distinct from workers. Univariate statistics showed that just a few characters for each colony were different among queens and workers (Table 2) in all colonies. Except for MTL in a worker-producing colony, all significant differences showed queens slightly larger than workers. Using the characters with significant differences in univariate statistics, covariance analysis (ANCOVA: Table 2) showed statistically signifi-

cant values in all these characters except for T1L in the pre-emergence colony, demonstrating that these characters differed in size and body proportions. ANCOVA was not tested in the absconding colony due to the presence of only one queen.

Discriminant function analysis (Table 3) showed also that only a few characters used in the model resulted in significant *p*-values. Wilk's Lambda values ranged from 0.64 to 0.8 (Table 10) in pre-emergence and worker-producing colonies and from 0.13 to 0.75 (one variable only) in

Table 2: Queen and worker means (in mm), Bonferroni t-test and ANCOVA analyses for each morphometrical variable in *Metapolybia docilis*.

Colony Phase	Characters	Means ± SD		Bonferroni t-test		ANCOVA (AL as covariate)			
		Queens	Workers	F	P>F	MS Effect	MS Error	F	p-level
Pre-emergence 9 queens and 64 workers	HL	2.00 ± 0.06	1.95 ± 0.07	1.89	0.06				
	HW	2.42 ± 0.06	2.41 ± 0.07	0.63	0.53				
	IDm	1.00 ± 0.04	0.99 ± 0.04	0.25	0.80				
	PW	1.86 ± 0.07	1.83 ± 0.06	1.79	0.08				
	MTL	0.48 ± 0.03	0.47 ± 0.03	0.72	0.48				
	AL	2.97 ± 0.12	2.90 ± 0.14	1.53	0.13				
	T2H*	2.02 ± 0.06	1.90 ± 0.10	3.19	0.002	0.05	0.01	7.91	0.01
	T1L*	2.22 ± 0.04	2.16 ± 0.08	2.20	0.03	0.01	0.003	2.41	0.13
	T2BW	0.50 ± 0.02	0.49 ± 0.04	0.77	0.45				
	WL	4.48 ± 0.10	4.40 ± 0.13	1.72	0.09				
Worker-producing 23 queens and 30 workers	HL	2.03 ± 0.04	2.02 ± 0.03	1.66	0.10				
	HW	2.44 ± 0.04	2.46 ± 0.05	-1.26	0.21				
	IDm	1.04 ± 0.02	1.04 ± 0.03	1.58	0.12				
	PW	1.28 ± 0.04	1.29 ± 0.07	-0.53	0.60				
	MTL*	0.50 ± 0.02	0.53 ± 0.03	-3.00	0.004	0.01	0.001	11.40	
	AL	3.25 ± 0.07	3.24 ± 0.09	0.60	0.55				
	T2H*	2.11 ± 0.08	1.96 ± 0.07	7.55	<0.001	0.31	0.005	62.74	
	T1L	2.25 ± 0.04	2.23 ± 0.06	1.44	0.16				
	T2BW	0.51 ± 0.02	0.50 ± 0.02	1.74	0.09				
	WL	4.56 ± 0.08	4.54 ± 0.10	0.62	0.54				
Male-producing 4 queens and 30 workers	HL	1.95 ± 0.05	2.00 ± 0.04	1.98	0.06				
	HW	2.43 ± 0.07	2.45 ± 0.03	0.82	0.42				
	IDm	1.03 ± 0.03	1.03 ± 0.03	0.12	0.90				
	PW*	1.42 ± 0.05	1.27 ± 0.04	-5.69	<0.001	0.08	0.002	41.39	<0.001
	MTL	0.54 ± 0.03	0.53 ± 0.03	-0.80	0.43				
	AL	3.06 ± 0.15	3.15 ± 0.09	1.73	0.09				
	T2H	1.98 ± 0.07	2.02 ± 0.11	0.60	0.55				
	T1L	2.20 ± 0.12	2.23 ± 0.05	1.02	0.32				
	T2BW	0.50 ± 0.00	0.48 ± 0.02	-0.85	0.40				
	WL	4.53 ± 0.16	4.55 ± 0.09	0.41	0.69				
Absconding 1 queen and 44 workers	HL	2.00	1.94 ± 0.05	1.10	0.28				
	HW	2.37	2.40 ± 0.05	-0.67	0.51				
	IDm	1.06	1.02 ± 0.05	0.87	0.39				
	PW	1.25	1.24 ± 0.08	0.05	0.96				
	MTL*	0.56	0.47 ± 0.04	2.19	0.04				
	AL	3.06	3.06 ± 0.11	0.02	0.99				
	T2H	2.00	1.95 ± 0.10	0.49	0.63				
	T1L	2.18	2.15 ± 0.07	0.39	0.70				
	T2BW	0.50	0.49 ± 0.03	0.13	0.90				
	WL	4.25	4.31 ± 0.11	-0.57	0.57				
Swarm 7 queens and 11 workers	HL	1.94 ± 0.04	1.92 ± 0.06	1.03	0.32				
	HW	2.37 ± 0.05	2.34 ± 0.08	0.87	0.40				

Table 2: Queen and worker means (in mm), Bonferroni t-test and ANCOVA analyses for each morphometrical variable in *Metapolybia docilis*. (Continued)

IDm*	1.01 ± 0.03	0.97 ± 0.03	2.69	0.02	0.01	0.001	9.03	0.01
PW	1.19 ± 0.04	1.18 ± 0.07	0.29	0.78				
MTL	0.50 ± 0.00	0.48 ± 0.03	1.18	0.26				
AL	3.00 ± 0.03	2.96 ± 0.03	0.50	0.63				
T2H*	1.94 ± 0.07	1.82 ± 0.08	3.28	0.005	0.06	0.01	9.71	0.01
T1L	2.15 ± 0.03	2.14 ± 0.09	0.11	0.91				
T2BW	0.48 ± 0.03	0.47 ± 0.03	0.69	0.50				
WL*	4.41 ± 0.07	4.26 ± 0.14	2.62	0.02	0.07	0.01	10.61	0.01

* Statistically significant

male-producing and swarm colonies. This indicates that in the former some individual characters were not good enough to discriminate castes, and caste differences could be determined only in association with these characters, which demonstrates low differentiation between castes (Fig. 2). In the latter, however, some characters appear to have a better power of discrimination (Fig. 2). Such differences suggest that variation during the colony cycle may occur as in other epiponines [11,17]. In fact, comparing actual groups with predicted groups through discriminant analysis (Table 3), it is evident that queens form more distinct groups in the later stages of the colony cycle. These results emphasize how queens, apart from being hardly distinguishable by size, exhibit some differences that are perhaps more related to shape than size.

Discussion

In *Metapolybia*, castes are hardly detectable by external morphology [1,28], and they are apparently determined by disputes among adults rather than by larval manipulation [6,30]. For this reason, young females have their ovaries suppressed in queen-right colonies, and only in orphanage young females develop their ovaries and become queens [6]. In our study, *Metapolybia docilis* exhibits slight differentiation between queens and workers, at least in the early stages of nest development, as compared to other taxa such as *Agelaia* spp. (*A. flavipennis* [31]; *A. areata*, [19]; *A. pallipes* and *A. multipicta* [22]; *A. vicina* [20,21]) that have a clear-cut caste differentiation. Also, the clear physiological distinction found between queen and worker ovaries (Fig. 2) suggest strong control of reproduction.

Group comparisons (Table 3) showed queens and workers as unique, well-defined groups, in the latter stages of the colony cycle. In addition, Wilks' Lambda values were relatively low for such colonies (Table 3), demonstrating that, in these colonies, castes can be relatively well-recognized by morphology in the later stages of the colony cycle. Such a pattern seems to be widespread for the Epiponini [11,17], and it is suggested that the increase in castes differences is probably due to the elimination of the

smaller queens. The fact that queens in the later stages of the colony cycle are slightly more distinct from those found in early stages suggest that the phenomenon of queen-selection [6] during the colony cycle may have a morphological component similar to other species previously studied [11,17]. In *Synoeca cyanea* [16], a similar pattern was identified. However, intermediate females were found in some colonies and differences between castes were always absent.

Differently from other species with increasing caste differences according to colony cycle, in *M. docilis*, size differences are very slight. Based on the plots of figure 2, castes are always intersecting each other, and in figure 2A, the growth rates, represented by different slopes detected by ANCOVA analyses (Table 2), are different between castes. Such results indicate that queens and workers differ more in shape than in size. Shape differences have been previously detected for *Apoica pallens* [24] but, in this case, it was accompanied by size differences, as in other species, such as *Epipona guerini* [18] and *Pseudopolybia difficilis* [32]. According to [24] it is possible that the first step of caste evolution in epiponines may be related to changes in body proportions rather than in size due to an ontogenetic reprogramming in the growth parameters (see also [33]). Thus, the almost absence of queen-worker size differences in *Metapolybia* illustrates how complex castes in epiponines is.

Low morphological differentiation accompanied by high physiological distinction was suggested for *Synoeca* and *Metapolybia* [6,30]. In these cases, only certain females that are already mated become queens. During cyclical oligogyny, a low number of queens would signal that new females would "take a chance" to become new queens. After queen repopulation, orphanage behaviors cease, and workers use aggressive behaviors to inhibit other new females from becoming queens. In this case, non-inseminated ovarian-developed females, common in the some epiponines [11] are completely suppressed during the entire colony cycle due to worker policing as suggested by [11]. In fact, queens' age (Fig. 1) reinforces such




Figure 2

Plots (in mm) of alitrunk length (AL) versus basal height tergum II (T₂H) in five colonies of *Metapolybia docilis*. Queens = black squares; Workers = white circles. Each ellipse encompasses 95% of the variation found in each group. A – discrimination between all queens and workers of *Metapolybia docilis* using the log-log plots.

Table 3: Discriminant morphometric variables between queens and workers in *Metapolybia docilis* based on discriminant function analyses using the stepwise procedure and classification results for group comparisons through discriminant analysis. Original data are based on mm.

Colony Phase	Variable	Step	Wilks Lambda	F	p	Observed classification	Predicted classification		Percent correct
Pre-emergence	T2H*	1	0.80	10.17	<0.001	Worker	61	3	95.31
	HW*	2	0.75	5.70	0.02	Queen	3	6	66.67
	PL	3	0.70	3.50	0.07	Total	64	9	91.7
	T2BW	4	0.69	3.53	0.06				
	TIL	5	0.66	2.88	0.09				
	MTL	6	0.67	2.70	0.10				
	IDm		0.66	2.98	0.09				
	HL		0.64	1.20	0.28				
	AL		0.64	1.16	0.29				
Worker-production	T2H*	1	0.80	57.00	<0.001		Worker	Queen	
	HW*	2	0.75	6.25	0.02	Worker	29	1	96.7
	HL	3	0.70	3.61	0.06	Queen	3	20	86.96
	MTL	4	0.69	3.03	0.09	Total	32	21	92.45
	IDX	5	0.66	1.71	0.20				
	TIL	6	0.67	2.40	0.13				
Male-production	PW*	1	0.75	32.34	<0.001	Worker	30	0	100
	TIL*	2	0.29	9.00	0.01	Queen	0	4	100
	HL	3	0.30	2.76	0.11	Total	30	4	100
	T2BW	4	0.32	2.74	0.11				
	AL		0.29	1.52	0.23				
	IDM		0.30	1.95	0.17				
	HW		0.29	1.24	0.28				
Swarm	T2H*	1	0.25	10.75	<0.001	Worker	11	0	100
	WL	2	0.37	3.66	0.08	Queen	0	7	100
	TIL*	3	0.18	8.43	0.01	Total	11	7	100
	PL	4	0.23	2.17	0.16				
	HL	5	0.24	3.92	0.07				
	HW*	6	0.19	5.40	0.04				
	IDm	7	0.13	1.01	0.34				

assumption, once pre-emergence colonies initiate with differently aged queens, and only older queens are found in worker-producing and male-producing colonies. In swarming colonies, the queen population is composed either of very young or very old individuals, suggesting that new reproductive are produced only during discrete windows of time.

The fact we found some distinction between castes would not reject the possibility of some young individuals (potential workers) may try to become queens [6,30]. Their options would be two-fold: they could lead the colony until a new queen repopulation occurs, or they could compete with those true queens during the early stages of the colony cycle. However, they would probably be removed from the colony as the cycle progresses [11].

Conclusions

As stated by [4] and [31], because epiponines frequently migrate, development of true castes would not be advantageous. This assertion may be correct in part, once castes are hard to detect. However, differently from the other "highly eusocial insects" (bees, ants, termites and vespine wasps) which bear pronounced size dimorphism, caste differences in epiponines evolved in such a way that imperceptible size differences are expected in most species, probably due to reprogramming of growth parameters [24]. The pattern of caste differences found in *Metapolybia docilis* supports this hypothesis.

Methods

Five colonies of *Metapolybia docilis* used in this work were collected at nightfall. They were collected into a plastic

bag containing ether-moistened cotton balls. Populations were fixed in Dietrich's solution and then kept in 70 % ethanol for dissection. In order to detect morphological differences between the castes, the following 10 external body parts were measured for each specimen under a binocular microscope; head width (HW), head length (HL), minimum interorbital distance (IDm), pronotum width (PW), mesoscutellum length (MTL), alitrunk length (AL), length of gastral tergite I (T1L), basal width of gastral tergite II (T2BW), height of T2 (T2H) and partial length of the forewing (WL) (see also Fig. 2 in [23] more details about these measurements). We examined ovarian conditions by dissecting all individuals under a stereomicroscope. In order to analyze insemination, the spermatheca was removed and put on a slide in a 1:1 solution of glycerin and alcohol (70%). The presence of sperm cells was detected by using a compound microscope. Numerical data were statistically analyzed in relation to the ovarian and spermathecal states.

Before statistical analysis, data were log transformed. Means and standard deviations were calculated from the ten morphological measurements. The Bonferroni *t* test was used for mean comparisons. The contribution of each variable to caste discrimination was examined using stepwise discriminant function analysis [34]. In this method, variables are successively added to the model based on the higher *F* to enter values; variable addition ceases when the *F*-ratio is no longer significant. Wilks' Lambda values were used to infer the individual contribution of each variable to the model. ANCOVA was used with alitrunk length (AL) as a covariate based on its efficiency in previous publications [18,24,32].

In order to assess the age of females, progressive pigmentation of the transverse apodeme was analyzed. The patterns used here were those that, according to [4,35] indicate a progressive increase in age. They were: LY (light yellow) for the youngest individuals, LB (light brown), DB (dark brown) and B (black) for the oldest individuals.

Authors' contributions

MVB participated in morphometric analyses, females' dissections. FBN and RZ participated in the design of the study, performed the statistical analysis and wrote the manuscript. All authors read and approved the final manuscript.

Acknowledgments

The authors acknowledge the financial support by FAPESP proc. 97/13204-9 and 01/02491-4. Special thanks to Sidnei Mateus for helping in field collections and Kurt M. Pickett (The Ohio State University) for corrections on the manuscript and important suggestions.

References

1. Richards OW: **The social wasps of the Americas excluding the Vespinae** London. *British Museum (Natural History)* 1978.
2. Carpenter JM: **Phylogenetic relationships and the origin of social behavior in the Vespidae** In: *The social biology of wasps* Edited by: Ross KG, Matthews RW. Cornell University Press, Ithaca, New York; 1991:7-32.
3. Jeanne RL: **The swarm-founding Vespidae** In: *The social biology of wasps* Edited by: Ross KG, Matthews RW. Cornell University Press, Ithaca, New York; 1991:191-231.
4. Richards OW and Richards MJ: **Observations on the social wasps of South America (Hymenoptera, Vespidae)** *Trans Roy Entomol Soc London* 1951, **102**:1-170.
5. Richards OW: **The biology of the social wasps (Hymenoptera, Vespidae)** *Biol Rev* 1971, **46**:483-528.
6. West-Eberhard MJ: **Temporary queens in *Metapolybia* wasps: non-reproductive helpers without altruism?** *Science* 1978, **200**:441-443.
7. Queller DC, Hughes CR and Strassmann JE: **Genetic relatedness in colonies of tropical wasps with multiple queens** *Science* 1988, **242**:1155-1157.
8. Queller DC, Negrón-Sotomayor JA, Hughes CR and Strassmann JE: **Queen number and genetic relatedness in a neotropical wasp, *Polybia occidentalis*** *Behav Ecol* 1993, **4**:7-13.
9. Hughes CR, Queller DC, Negrón-Sotomayor JA, Strassmann JE, Solis C and Gastreich KR: **The maintenance of high genetic relatedness in multi-queen colonies of social wasps** In: *Queen number and sociality in insects* Edited by: Keller L. Oxford University press, New York; 1993.
10. Strassmann JE, Goodnight KF, Kihner CJ and Queller DC: **The genetic structure of swarms and the timing of their production in the queen cycles of neotropical wasps** *Molecular Ecology* 1998, **7**:709-718.
11. Noll FB and Zucchi R: **Caste and influence of the colony cycle in swarm-founding polistine wasps (Hymenoptera, Vespidae, Epiponini)** *Insectes Sociaux* 2002, **49**:62-74.
12. Mateus S, Noll FB and Zucchi R: **Morphological caste differences in neotropical swarm-founding polistine wasps: *Parachartergus smithii* (Hymenoptera: Vespidae)** *J New York Entomol Soc* 1997, **105**:129-139.
13. Shima SN, Noll FB, Zucchi R and Yamane S: **Morphological caste differences in the Neotropical swarm-founding polistine Wasps IV. *Pseudopolybia vespiceps*, with preliminary considerations on the role of intermediate females in social organization of the Epiponini (Hymenoptera, Vespidae)** *J Hym Res* 1998, **7**:280-295.
14. Mateus S, Noll FB and Zucchi R: **Caste differences and related bionomic aspects of *Chartergellus communis*, a neotropical swarm-founding polistine wasp (Hymenoptera: Vespidae: Polistinae: Epiponini)** *J New York Entomol Soc* 1999, **107**:390-405.
15. Shima SN, Noll FB and Zucchi R: **Morphological caste differences in the neotropical swarm-founding polistine wasp, *Brachygastra lecheguana* (Hymenoptera: Vespidae, Polistinae, Epiponini)** *Sociobiology* 2000, **36**:41-52.
16. Noda SCM, Shima SN and Noll FB: **Morphological and physiological caste differences in *Synoeca cyanea* (Hymenoptera, Vespidae, Epiponini) according to the ontogenetic development of the colonies** *Sociobiology* 2003, **41**:547-570.
17. Noll FB and Zucchi R: **Increasing caste differences related to life cycle progression in some neotropical swarm-founding polygynic wasps (Hymenoptera: Vespidae: Epiponini)** *Ethol Ecol Evol* 2000, **12**:43-65.
18. Hunt JH, Schmidt DK, Mulkey SS and Williams MA: **Caste dimorphism in *Epipona guerini* (Hymenoptera: Vespidae): Further evidence for larval determination** *J Kansas Entomol Soc* 1996, **69**:362-369.
19. Jeanne RL and Fagen R: **Polymorphism in *Stelopolybia areata* (Hymenoptera, Vespidae)** *Psyche* 1974, **81**:155-166.
20. Sakagami SF, Zucchi R, Yamane S, Noll FB and Camargo JMF: **Morphological caste differences in *Agelaia vicina*, the neotropical swarm-founding Polistinae wasp with the largest colony size among social wasps (Hymenoptera, Vespidae)** *Sociobiology* 1996, **28**:207-223.
21. Baio MV, Noll FB, Zucchi R and Simões D: **Non-allometric caste differences in *Agelaia vicina* (Hymenoptera, Vespidae, Epiponini)** *Sociobiology* 1998, **32**:465-476.

22. Noll FB, Simões D and Zucchi R: **Morphological caste differences in the neotropical swarm-founding Polistine wasps: *Agelaia m. multipicta* and *A. p. pallipes* (Hymenoptera Vespidae)** *Ethology Ecology & Evolution* 1997, **9**:361-372.
23. Shima SN, Yamane S and Zucchi R: **Morphological Caste Differences in Some Neotropical Swarm-founding Polistine Wasps I. *Apoica flavissima* (Hymenoptera, Vespidae)** *Jpn J Ent* 1994, **62**:811-822.
24. Jeanne RL, Graf CA and Yandell BS: **Non-Size-Based morphological castes in a social insect** *Naturwissenschaften* 1995, **82**:296-298.
25. Shima SN, Yamane S and Zucchi R: **Morphological Caste Differences in Some Neotropical Swarm-founding Polistine Wasps II. *Polybia dimidiata* (Hymenoptera, Vespidae)** *Jpn J Ent* 1995, **64**:131-144.
26. Smethurst ME and Carpenter JM: **A new species of *Metapolybia ducke* from Central America (Hymenoptera: Vespidae; Polistinae)** *J New York Ent Soc* 1998, **105**:180-185.
27. Itô Y, Yamauchi K and Tsuchida K: **Reproductive condition of females in some swarm-founding wasps in Panama, as compared with some independent-founding species (Hymenoptera: Vespidae)** *Sociobiology* 1997, **29**:269-276.
28. Carpenter JM and Ross KG: **Colony composition in four species of Polistinae from Suriname, with a description of the larva of *Brachygastra scutellaris* (Hymenoptera, Vespidae)** *Psyche* 1984, **91**:237-250.
29. Karsai I and Wenzel JW: **Organization and regulation of nest construction behavior in *Metapolybia* wasps** *J Ins Behav* 2000, **13**:111-140.
30. West-Eberhard MJ: **Intragroup selection and evolution of insect societies** *In Natural selection and social behavior* Edited by: Alexander RD, Tinkle DW. Chiron Press, New York; 2000:3-17.
31. Evans H and West-Eberhard MJ: **The wasps** *Univ Michigan, Ann Arbor* 1970.
32. Jeanne RL: **Non-allometric queen-worker dimorphism in *Pseudopolybia difficilis* (Hymenoptera: Vespidae)** *J Kansas Entomol Soc* 1996, **69**:370-374.
33. Wheeler DE: **The development basis of worker castes polymorphism in ants** *Am Nat* 1991, **138**:1218-1238.
34. Rao CR: **Linear statistical inference** *John Wiley and Sons, New York* 1973.
35. Forsyth A: **Studies on the behavioral ecology of polygynous social wasps** *Doctoral Dissertation, Harvard University, Cambridge* 1978.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

